

COLÁISTE NA TRÍONÓIDE, BAILE ÁTHA CLIATH | TRINITY COLLEGE DUBLIN
Ollscoil Átha Cliath | The University of Dublin

JF Chemistry 1101 2010 Introduction to Physical Chemistry: Acid Base and Solution Equilibria.




Dr Mike Lyons
School of Chemistry
melyons@tcd.ie



Required Reading Material.

- Silberberg, Chemistry, 4th edition.
 - Chapter 18.
 - Acid/base equilibria. pp.766-813.
 - Chapter 19.
 - Ionic equilibria in aqueous systems. pp.814-862.
- Kotz, Treichel and Weaver, 7th edition.
 - Chapter 17 (Chemistry of Acids and Bases) & Chapter 18 (Principles of reactivity: other aspects of ionic equilibria), pp.760-859.
- Chemistry³, Burrows et al.
 - Chapter 6, Acids & bases, pp.263-299.



Review : Kotz Chapter 3 for simple acid/base definitions.


Lecture 13.

Acid/base chemistry :
Simple ideas: Arrhenius,
Bronsted-Lowry, Lewis.






Kotz: section 3.7, pp.131-139.
Section 17.1, pp.761-762.

Acid and Bases



▲ The tartness of lemons and oranges comes from the weak acid citric acid. The acid is found widely in nature and in many consumer products.
(Charles D. Winters)




▲ The sting of ants is due to the weak acid formic acid, HCO_2H .


(Gallo Images/@ CORBIS)



Acid and Bases






Acid and Bases




Arrhenius (or Classical) Acid-Base Definition


- An **acid** is a neutral substance that contains hydrogen and dissociates or ionizes in water to yield hydrated protons or hydronium ions H_3O^+ .
- A **base** is a neutral substance that contains the hydroxyl group and dissociates in water to yield hydrated hydroxide ions OH^- .
- Neutralization** is the reaction of an H^+ (H_3O^+) ion from the acid and the OH^- ion from the base to form water, H_2O .
- These definitions although correct are limited in that they are not very general and do not
- Give a comprehensive idea of what acidity and basicity entails.



Arrhenius acid is a substance that produces H^+ (H_3O^+) in water.




Arrhenius base is a substance that produces OH^- in water.



Acids and bases: Bronsted/Lowry definition.

- **Bronsted/Lowry Acid (HA):**
 - An acid is a species which **donates** a proton
- **Bronsted/Lowry Base (B):**
 - A base is a species which **accepts** a proton.
- These definitions are quite general and refer to the reaction between an acid and a base.
- An acid must contain H in its formula; HNO_3 and H_2PO_4^- are two examples, all Arrhenius acids are Bronsted-Lowry acids.
- A base must contain a lone pair of electrons to bind the H^+ ion; a few examples are NH_3 , CO_3^{2-} , F^- , as well as OH^- . Bronsted-Lowry bases are not Arrhenius bases, but all Arrhenius bases contain the Bronsted-Lowry base OH^- .




- In the Bronsted-Lowry perspective:
one species donates a proton and another species accepts it: an acid-base reaction is a proton transfer process.

Chemistry³ section 6.1. pp.264-267.


Kotz 7th ed. Section 17.1. pp.761-765

BL acid/base equilibria.




- Proton donation and acceptance are **dynamic** processes for all acids and bases. Hence a **proton transfer equilibrium** is rapidly established in solution.
- The equilibrium reaction is described in terms of conjugate acid/base pairs.
- The conjugate base (CB) of a BL acid is the base which forms when the acid has donated a proton.
- The conjugate acid (CA) of a BL base is the acid which forms when the base has accepted a proton.
- A conjugate acid has one more proton than the base has, and a conjugate base one less proton than the acid has.
- If the acid of a conjugate acid/base pair is strong (good tendency to donate a proton) then the conjugate base will be weak (small tendency to accept a proton) and vice versa.

Acid : proton donor
Base : proton acceptor



A Brønsted **acid** is a proton donor

A Brønsted **base** is a proton acceptor



15.1

Brønsted-Lowry Acid-Base Definition

An acid is a proton donor, any species which donates a H^+ .

A base is a proton acceptor, any species which accepts a H^+ .

An acid-base reaction can now be viewed from the standpoint of the reactants AND the products.

An acid reactant will produce a base product and the two will constitute an acid-base conjugate pair.

Table 18.4 The Conjugate Pairs in Some Acid-Base Reactions

Conjugate Pair

Acid + Base \rightleftharpoons Base + Acid

Conjugate Pair

Reaction 1	HF	+	H ₂ O	\rightleftharpoons	F ⁻	+	H ₃ O ⁺
Reaction 2	HCOOH	+	CN ⁻	\rightleftharpoons	HCOO ⁻	+	HCN
Reaction 3	NH ₄ ⁺	+	CO ₃ ²⁻	\rightleftharpoons	NH ₃	+	HCO ₃ ⁻
Reaction 4	H ₂ PO ₄ ⁻	+	OH ⁻	\rightleftharpoons	HPO ₄ ²⁻	+	H ₂ O
Reaction 5	H ₂ SO ₄	+	N ₂ H ₅ ⁺	\rightleftharpoons	HSO ₄ ⁻	+	N ₂ H ₆ ²⁺
Reaction 6	HPO ₄ ²⁻	+	SO ₃ ²⁻	\rightleftharpoons	PO ₄ ³⁻	+	HSO ₃ ⁻

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Table 15.2 Relative Strengths of Conjugate Acid-Base Pairs

	Acid	Conjugate Base	
<div style="display: flex; flex-direction: column; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">Acid strength increases</div> <div style="margin-top: 20px;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">Weak acids</div> </div> </div>	HClO ₄ (perchloric acid)	ClO ₄ ⁻ (perchlorate ion)	<div style="display: flex; flex-direction: column; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">Base strength increases</div> </div>
	HI (hydroiodic acid)	I ⁻ (iodide ion)	
	HBr (hydrobromic acid)	Br ⁻ (bromide ion)	
	HCl (hydrochloric acid)	Cl ⁻ (chloride ion)	
	H ₂ SO ₄ (sulfuric acid)	HSO ₄ ⁻ (hydrogen sulfate ion)	
	HNO ₃ (nitric acid)	NO ₃ ⁻ (nitrate ion)	
	H ₃ O ⁺ (hydronium ion)	H ₂ O (water)	
	HSO ₄ ⁻ (hydrogen sulfate ion)	SO ₄ ²⁻ (sulfate ion)	
	HF (hydrofluoric acid)	F ⁻ (fluoride ion)	
	HNO ₂ (nitrous acid)	NO ₂ ⁻ (nitrite ion)	
	HCOOH (formic acid)	HCOO ⁻ (formate ion)	
	CH ₃ COOH (acetic acid)	CH ₃ COO ⁻ (acetate ion)	
	NH ₄ ⁺ (ammonium ion)	NH ₃ (ammonia)	
	HCN (hydrocyanic acid)	CN ⁻ (cyanide ion)	
	H ₂ O (water)	OH ⁻ (hydroxide ion)	
	NH ₃ (ammonia)	NH ₂ ⁻ (amide ion)	



15.4

Table 6.1 The relative strengths of acids and their conjugate bases



	Conjugate acid		Conjugate base	
Strongest acid	$\text{HClO}_4(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{ClO}_4^-(\text{aq})$	Weakest base
	perchloric acid		perchlorate ion	
	$\text{HCl}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{Cl}^-(\text{aq})$	
	hydrochloric acid		chloride ion	
	$\text{H}_2\text{SO}_4(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{HSO}_4^-(\text{aq})$	
	sulfuric acid		hydrogensulfate ion	
	$\text{H}_3\text{O}^+(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{H}_2\text{O}(\text{l})$	
	oxonium ion		water	
	$\text{CH}_3\text{CO}_2\text{H}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{CH}_3\text{CO}_2^-(\text{aq})$	
	ethanoic acid		ethanoate ion	
	$\text{NH}_4^+(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{NH}_3(\text{aq})$	
	ammonium ion		ammonia	
	$\text{H}_2\text{O}(\text{l}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{OH}^-(\text{aq})$	
	water		hydroxide ion	
Weakest acid	$\text{C}_2\text{H}_5\text{OH}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons$		$\text{H}_3\text{O}^+(\text{aq}) + \text{C}_2\text{H}_5\text{O}^-(\text{aq})$	Strongest base
	ethanol		ethoxide ion	

SAMPLE PROBLEM 18.4: Identifying Conjugate Acid-Base Pairs

PROBLEM: The following reactions are important environmental processes. Identify the conjugate acid-base pairs.




PLAN: Identify proton donors (acids) and proton acceptors (bases).



Strong and weak acids.

- Strong acids dissociate completely into ions in water:




In a dilute solution of a strong acid, almost no HA molecules exist: $[H_3O^+] = [HA]_{init}$ or $[HA]_{eq} = 0$

$$Q_c = \frac{[H_3O^+][A^-]}{[HA][H_2O]} \text{ at equilibrium, } Q_c = K_c \gg 1$$

Nitric acid is an example: $HNO_3(l) + H_2O(l) \longrightarrow H_3O^+_{(aq)} + NO_3^-_{(aq)}$

- Weak acids dissociate very slightly into ions in water:




In a dilute solution of a weak acid, the great majority of HA molecules are undissociated: $[H_3O^+] \ll [HA]_{init}$ or $[HA]_{eq} = [HA]_{init}$

$$Q_c = \frac{[H_3O^+][A^-]}{[HA][H_2O]} \text{ at equilibrium, } Q_c = K_c \ll 1$$



The Extent of Dissociation for Strong and Weak Acids

Key concept :
Acid/base strength quantified in terms of **extent** or **degree** of **dissociation**.
An acid or base is classified as **strong** if it is **fully ionized** in solution (e.g. HCl, NaOH).
An acid or base is classified as **weak** if only a **small fraction** is ionized in solution (e.g. CH_3COOH , NH_3).





Extent of dissociation : strong acid.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Extent of dissociation: weak acid.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Reactivity of strong and weak acids.

1M HCl(aq)



1M $\text{CH}_3\text{COOH(aq)}$

Classifying the Relative Strengths of Acids.

Strong acids.

There are two types of strong acids:

- The hydrohalic acids HCl , HBr , and HI
- Oxoacids in which the number of O atoms exceeds the number of ionizable H atoms by two or more, such as HNO_3 , H_2SO_4 , HClO_4

Weak acids.

There are many more weak acids than strong ones. Four types, with examples, are:

- The hydrohalic acid HF
- Those acids in which H is bonded to O or to halogen, such as HCN and H_2S
- Oxoacids in which the number of O atoms equals or exceeds by one the number of ionizable H atoms, such as HClO , HNO_2 , and H_3PO_4
- Organic acids (general formula RCOOH), such as CH_3COOH and $\text{C}_6\text{H}_5\text{COOH}$.

Classifying the Relative Strengths of Bases.

Strong bases.

- Soluble compounds containing O^{2-} or OH^- ions are strong bases. The cations are usually those of the most active metals: M_2O or MOH , where M = Group 1A(1) metals (Li, Na, K, Rb, Cs).
- MO or $M(OH)_2$, where M = Group 2A(2) metals (Ca, Sr, Ba) [MgO and $Mg(OH)_2$ are only slightly soluble, but the soluble portion dissociates completely.]


Weak bases.

- Many compounds with an electron-rich nitrogen are weak bases (none are Arrhenius bases). The common structural feature is an N atom that has a lone electron pair in its Lewis structure.
 - Ammonia (NH_3)
 - Amines (general formula RNH_2 , R_2NH , R_3N), such as $CH_3CH_2NH_2$, $(CH_3)_2NH$, $(C_3H_7)_3N$, and C_5H_5N




Strong Electrolyte - 100% dissociation



Weak Electrolyte - not completely dissociated






Strong Acids are strong electrolytes





15.4

Weak Acids are weak electrolytes




Strong Bases are strong electrolytes



15.4

Weak Bases are weak electrolytes



Conjugate acid-base pairs:


- The conjugate base of a strong acid has no measurable strength.
- H_3O^+ is the strongest acid that can exist in aqueous solution.
- The OH^- ion is the strongest base that can exist in aqueous solution.

15.4

Representing Protons





- Both representations of the proton are equivalent.
- H_5O_2^+ (aq), H_7O_3^+ (aq), H_9O_4^+ (aq) have been observed.
- We will use $\text{H}^+(\text{aq})$!

The Nature of the Hydrated Proton



The hydrated proton is quite a complex entity. It is usually represented in shorthand form as $\text{H}^+(\text{aq})$. A better representation is in terms of the hydronium ion H_3O^+ . We will adopt this representation a lot. The real situation is more complex. The H_3O^+ ion binds to other water molecules forming a mixture of species with the general formula $\text{H}(\text{H}_2\text{O})_n^+$. In fact the structural details of liquid water is still a hot item of research.

What is H^+ (aq)?



Yet more sophistication: Lewis acidity


An **Arrhenius acid** is defined as a substance that produces H^+ (H_3O^+) in water.

A **Brønsted acid** is defined as a proton donor


A **Lewis acid** is defined as a substance that can accept a pair of electrons.

A **Lewis base** is defined as a substance that can donate a pair of electrons

G.N. Lewis 1875-1946



See Kotz section 17.9
pp.789-798.




http://en.wikipedia.org/wiki/Gilbert_N._Lewis

Electron-Pair Donation and the Lewis Acid-Base Definition

The Lewis acid-base definition :

- A **base** is any species that *donates* an electron pair.
- An **acid** is any species that *accepts* an electron pair.


Protons act as Lewis acids in that they accept an electron pair in all reactions:




The product of any Lewis acid-base reaction is called an **adduct**, a single species that contains a new covalent bond.

- A Lewis base has a lone pair of electrons to donate.
- A Lewis acid has a vacant orbital

Lewis Acid/Base Reaction



Lewis Acids and Bases





No protons donated or accepted!

15.12


Reaction of NH_3 with $\text{Cu}^{2+}(\text{aq})$



PLAY MOVIE



Copper-ammonia coordinate covalent bond.




Ammonia



Water

PLAY MOVIE




The Lewis
Acid-Base
Chemistry
of
Nickel(II)

Lewis Acids & Bases





+ DMG

DMG =
dimethylglyoxime, a
standard reagent
to detect
nickel(II)



Lewis Acid-Base Interactions in Biology




Heme group

- The heme group in hemoglobin can interact with O_2 and CO .
- The Fe ion in hemoglobin is a Lewis acid
- O_2 and CO can act as Lewis bases


Lewis Acids & Bases

Many complex ions containing water undergo **HYDROLYSIS** to give acidic solutions.




Lewis Acids & Bases

This explains why water solutions of Fe^{3+} , Al^{3+} , Cu^{2+} , Pb^{2+} , etc. are acidic.



Amphoterism of $\text{Al}(\text{OH})_3$



(a) Add $\text{NH}_3(\text{aq})$

Adding aqueous ammonia to a soluble salt of Al^{3+} leads to a precipitate of $\text{Al}(\text{OH})_3$.

(b) Add $\text{NaOH}(\text{aq})$

Adding a strong base (NaOH) to $\text{Al}(\text{OH})_3$ dissolves the precipitate. Here, aluminum hydroxide acts as a Lewis acid toward the Lewis base OH^- and forms the soluble sodium salt of the complex ion $[\text{Al}(\text{OH})_4]^-$.


(c) Add $\text{HCl}(\text{aq})$

$\text{Al}(\text{OH})_3$ dissolves when a strong acid (HCl) is added. In this case, $\text{Al}(\text{OH})_3$ acts as a Brønsted base and forms a soluble aluminum salt and water.

© Brooks/Cole, Cengage Learning

Lewis Acids & Bases

This explains **AMPHOTERIC** nature of some metal hydroxides.



Here $\text{Al}(\text{OH})_3$ is a Brønsted base.



Here $\text{Al}(\text{OH})_3$ is a Lewis acid.

